



Which primary vaccine series can I take?



Ages 6 months - 4 years

Two doses
3-8 weeks apart, then a third
dose at least 8 weeks later.

Ages 5 years+

Two doses
3 - 8 weeks apart

Anyone with an
immunocompromising condition
should receive a third dose



Ages 6 months +

Two doses
4-8 weeks apart



Ages 12 years +

Two doses
3-8 weeks apart

People with moderately or severely compromised immune systems, allergies, or received the Johnson & Johnson/Janssen COVID-19 shot may follow a different primary vaccine schedule. Learn more from a clinician or certified Community Health Worker at helpguidethrive.org



Who should get a booster:

Ages 5 +

When to get a booster:

At least 2 months after the
second primary dose.

Anyone 6 months - 4 years old who
completed the three-dose primary series
is ineligible for a booster at this time.

C.D.C. recommends you get:

Ages 6 months to 4 years:

Moderna bivalent

Age 5 years:

Pfizer-BioNTech bivalent

Ages 6 years +:

Pfizer-BioNTech bivalent or
Moderna bivalent



Who should get a booster:

Ages 6 months +

When to get a booster:

At least 2 months after
completing a primary
vaccine series

C.D.C. recommends you get:

Ages 6 months to 4 years old:

Moderna bivalent

Ages 5 years +:

Pfizer-BioNTech bivalent
or Moderna bivalent



Who should get a booster:

Ages 12 years +

When to get a booster:

At least two months

At least 2 months after completing
a primary vaccine series

C.D.C. recommends you get:

Ages 12-17:

Pfizer-BioNTech bivalent
or Moderna bivalent



Which booster should I take?